Supernumerary Tooth: A Common Odontostomatologic Anomaly Of Number Of Teeth

Muhammad Shahrukh Khan Sadiq, Daud Mirza

INTRODUCTION:

Supernumerary tooth (ST) is a developmental disturbance of number of teeth characterized by teeth present in excess of normal dental formula of deciduous or permanent dentition.^{1,2,3,4} It may occur in either dental arch, maxillary or mandibular.⁵ Literature has shown marked predilection in maxilla over mandible.^{6,7} The first documented report of supernumerary teeth has been revealed in human fossils that are approximately 11,000 years old.⁸

Keywords: Supernumerary teeth, Hyperdontia, Mesiodens, Paramolars, Distomolars

METHODOLOGY:

Search engine of Google was utilized with various keywords and phrases to search articles related to supernumerary teeth from 2000-2018. Key words and phrases as supernumerary teeth, history of supernumerary teeth, etiology of supernumerary teeth, prevalence of supernumerary teeth, types of supernumerary teeth, diagnosis of supernumerary teeth, complications of supernumerary teeth and management of supernumerary teeth etc. were used. A total of 45 articles including reviews, original articles, case reports and textbooks were selected.

LITERATURE REVIEW:

Etiology:

The exact etiology of ST is unknown but, three main theories are proposed by researchers, in addition to the combination of genetic and environmental factors:

- The first theory is of phylogenetic reversion, which states that ST may be an atavistic appearance of fourth molar of primitive dental formula and signifies evolution.
- Second is dichotomy theory, which focuses on third tooth bud arising from dental lamina near permanent tooth bud or by splitting of permanent bud itself.
- Finally, there is theory of hyperactivity which is considered to be the most widely accepted. According to this theory, ST appear because of the independent hyperactivity of dental lamina that is localized in nature. Rudimentary form of tooth arises from proliferation of epithelial remnants of dental lamina induced by presence of complete dentition.^{9,10}

Prevalence:

Muhammad Shahrukh Khan Sadiq
Lecturer, Oral Pathology, BUMDC

Daud Mirza
Associate Professor and Head of Department Oral
Pathology, (BUMDC). Email:

Received: 20-02-18
Revised: 28-02-18
Accepted: 10-03-18

a. At International level:

The prevalence of ST varies around the world, study conducted in Spanish population showed its prevalence from 0.3% to 3.8% in their study subjects.¹¹ A Hungarian study conducted by Gabris on 2,219 patients reported prevalence of 1.53% in indigenous subjects.¹² Nayak and colleagues reported the prevalence of ST was 0.09-0.29% among Indian population.¹³ No significant sex distribution is seen in the primary dentition but males are affected approximately twice as compared to females in permanent teeth.^{14,15,16,17}

b. At National level:

In Pakistan, few extensive researches have been presented on the topic of ST. Prevalence of ST in orthodontic patients was 3.9% as reported by Farhat Amin at University of Lahore. Mesiodens were present in 0.87%, paramolars in 2.6% and distomolar was present in only in 0.43%. Gender wise, females were more affected than males by the ratio of 2:1 and left to right side ratio was found to be 5:2.^{16,17,18}

Diagnosis:

The ratio of ST in terms of impacted to erupted is 3:1.¹⁹ They are mostly diagnosed by clinical examination and routine radiographs of oral cavity. Often, clinical signs such as wide diastema, ectopic or eruption failure of permanent tooth, persistence of deciduous dentition and obvious visual presence of additional tooth aid the clinician in its diagnosis. Detailed history taking, proper clinical examination, thorough investigation, early diagnosis and appropriate treatment plan is essential for its management.^{20, 21}

Classification:

A range of classification standards are used to categorize ST.²² According to chronology, ST are classified as predeciduous, past permanent, or complementary. On the basis of morphology they can be odontoma, supplemental or eumorphic and rudimentary or dysmorphic (subdivided into conical, tuberculate, and molariform types) and according to orientation they can

be vertical, inverted, and transverse.^{24,25} On the basis of location, supernumerary teeth that occur between or posterior to the maxillary central incisors are referred as "mesiodens".^{26,27} Supernumerary structures found in molar region can be grouped as paramolars and distomolars. Distomolars are located distal or distolingually to wisdom molars.²⁸ Paramolars are usually smaller in size and dysmorphic in form, being located either buccally or lingually to one of the molar teeth.⁵ Mostly the occurrence of paramolars is unilateral and they are rarely observed bilaterally.²⁹

Complications:

Numerous complications accompany ST. Some of these complications can be crowding, malocclusion, delayed eruptions, localized periodontitis, caries, root resorption, dentigerous cyst and abnormal root development.^{30,31,32} Not only they are a source of complications but also interfere in alveolar bone grafting and implant placement.³³ According to the report by Parolia and Kundabala, adjacent molar is most likely to be affected with caries in case of maxillary paramolars, that are bilateral in nature.⁵

Association with syndromes:

ST are often associated with syndromes. Some of these are Cleidocranial Dysplasia, Gardner's syndrome, Fabry-Anderson syndrome, Ellis-van Creveld syndrome, cleft lip and palate. Less common associated syndromes involve Gorlin-Chaudhry-Moss syndrome, Willaim's syndrome, Hallermann-Streiff syndrome, orofaciogigital syndrome etc.^{34,35}

Management:

The management of ST involves endodontic therapy, retention of the additional tooth or extraction. The removal of ST is advised when:

- There is associated pathology
- Permanent tooth eruption has been delayed due to the presence of ST
- Increased risk of caries due to the presence of ST which makes the area inaccessible to maintain oral hygiene
- Altered eruption or displacement of adjacent tooth is evident
- There are severely rotated teeth leading to further complication
- Orthodontic treatment needs to be carried out to align the teeth
- Its presence would compromise alveolar bone grafting and implant placement
- There is compromised esthetic and functional status.^{36,37,38}

Conclusion:

ST is a pervasive developmental anomaly among the world

populace. The etiology of this anomaly still remains unclear. Various tooth complications are part and parcel of supernumerary teeth therefore, on diagnosis each case should be managed appropriately in order to mitigate the aesthetic and functional compromise to the dentition. The condition demands an efficient and effective treatment by the clinician.

Figure 1: Intraoral view showing mesiodens²⁵

Figure 2: Intraoral view showing fractured maxillary left permanent central incisor and mesiodens (arrow)²⁶

REFERENCES:

- R. Fuentes, E. Borie, and V. Beltrán, "Radiographical and macroscopical visualization of fourth molars: a report of four cases in maxilla," *International Journal of Morphology*, vol. 30, no. 1, pp. 115–118, 2012.
- R. S. Omer, R. P. Anthonappa, and N. M. King, "Determination of the optimum time for surgical removal of unerupted anterior supernumerary teeth," *Pediatric Dentistry*, vol. 32, no. 1, pp. 14–20, 2010.
- Chaudhary, S., Chaitra, T.R., Sultan, S., Arora, R. Supernumerary teeth in primary dentition. *BMJ Case Rep*. 2013.
- Celikoglu M, Kamak H, Oktay H. Prevalence and characteristics of supernumerary teeth in a non-syndrome Turkish population: Associated pathologies and proposed treatment. *Med Oral Patol Oral Cir Bucal*. 2010;15:e575–8.
- Parolia and M. Kundabala, "Bilateral Maxillary paramolars and endodontic therapy: a rare case report," *Journal of Dentistry Tehran University Medical Sciences*, vol. 7, no. 2, pp. 107–111, 2010.
- P. M. Omal, V. Jacob, J. Lonapan, and A. Kurian, "Bilateral fourth molars with paramolars in maxilla," *Kerala Dental Journal*, vol. 34, no. 2, pp. 277–279, 2011.
- N. B. Nagaveni, K. V. Umashankar, N. B. Radhika, P. B. Reddy,

- and S. Manjunath, "Maxillary paramolars report of a case and literature review," *Archives of Orofacial Sciences*, vol. 5, no. 1, pp. 24–28, 2010.
8. Benazzi S, Buti L, Franzo L, Kullmer O, Winzen O, Gruppioni G. Report of three fused primary human teeth in an archaeological material. *Int J Osteoarchaeol*. 2009 doi: 10.1002/oa.1056.
9. R. P. Anthonappa, R. S. M. Omer, and N. M. King, Characteristics of 283 supernumerary teeth in Southern Chinese children, *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontology*, vol. 105, no. 6, pp. e48–e54, 2008.
10. Xi Lu, Fang Yu, Junjun Liu, Wenping Cai, Yumei Zhao, Shouliang Zhao & Shangfeng Liu, The epidemiology of supernumerary teeth and the associated molecular mechanism, *Organogenesis*, 2017 13:3, 71–82.
11. Ferrés-Padró E, Prats-Armengol J, Ferrés-Amat E. A descriptive study of 113 unerupted supernumerary teeth in 79 pediatric patients in Barcelona. *Med Oral Patol Oral Cir Bucal*. 2009;14:E146–52.
12. Gabris K, Fabian G, Kaan M, Rozsa N, Tarjan I. Prevalence of hypodontia and hyperdontia in paedodontic and orthodontic patients in Budapest. *Community Dent Health* 2006; 23: 80–82.
13. G. Nayak, S. Shetty, S. Inderpreet, and D. Pitalia, "Paramolars—a supernumerary molar: a case report and over view," *Dental Research Journal*, vol. 9, pp. 797–803, 2012.
14. Menardia -Pejuan V. Supernumerary molars. A review of 53 cases. *It Res Sci stomatology*. 2000;402(2–3):101–5.
15. Kinirons MJ. Unerupted premaxillary supernumerary teeth. A study of their occurrence in males and females. *Br Dent J*. 1982;153:110.
16. E. C. Kuchler, A. G. Costa, C. Costa Mde, A. R. Vieira, and J. M. Granjeiro, "Supernumerary vary depending on gender," *Brazilian Oral Research*, vol. 25, pp. 76–79, 2011.
17. D. Rajab and M. A. M. Hamdan, "Supernumerary teeth: review of the literature and a survey of 152 cases," *International Journal of Paediatric Dentistry*, vol. 12, no. 4, pp. 244–254, 2002.
18. Amin F. Prevalence of hypodontia in orthodontic patients in a Pakistani sample-a study. *Pak Oral Dent J* 2010; 30: 142–145.
19. Gupta S, Marwah N. Impacted Supernumerary Teeth—Early or Delayed Intervention: Decision Making Dilemma? *International Journal of Clinical Pediatric Dentistry*. 2012;5(3):226–230. doi:10.5005/jp-journals-10005-1173.
20. Garvey MT, Barry HJ, Blake M. Supernumerary teeth- an overview of classification, diagnosis and management. *J Can Dent Assoc*. 1999;65:612–6.
21. R. P. Anthonappa, N. M. King, and A. B. M. Rabie, "Diagnostic tools used to predict the prevalence of supernumerary teeth: a meta-analysis," *Dentomaxillofacial Radiology*, vol. 41, no. 6, pp. 444–449, 2012.
22. Cakan DG, Ulkur F, Taner T. The genetic basis of dental anomalies and its relation to orthodontics. *Eur J Dent*. 2013;7:S143–7.
23. Ramesh K, Venkataraghavan K, Kunjappan S, Ramesh M. Mesiodens: A clinical and radiographic study of 82 teeth in 55 children below 14 years. *Journal of Pharmacy & Bioallied Sciences*. 2013;5(Suppl 1):S60–S62. doi:10.4103/0975-7406.113298.
24. Arandi NZ, Distomolars; an overview and 3case reports. *Dent Oral Craniofac Res* 4: DOI: 10.15761/DOCR.1000236, 2017.
25. Ozkan A, Dag H, Altug HA, Sencimen M. Bilateral Double Maxillary Paramolars: A Rare Case Report. *Journal of Clinical and Diagnostic Research? : JCDR*. 2017;11(8):ZD04-ZD05.
26. Gorlin RJ, Cohen MM, Hennekam RC. *Syndromes of the head and neck*. 4th ed. Oxford: Oxford University Press; 2001. p. 547.p. 1108.
27. Mukhopadhyay, S. Mesiodens: a clinical and radiographic study in children. *J Indian Soc Pedod Prev Dent*. 2011;29:34–38.
28. Victor Beltrán, Mario Cantín, Eduardo Borie, Ramón Fuentes, and Wilfried Engelke, "Endoscopic Removal of a Supernumerary Premolar in the Mandible during a Dental Implant Placement," *Case Reports in Dentistry*, vol. 2014, Article ID 975470, 5 pages, 2014. doi:10.1155/2014/975470.
29. Rajab LD, Hamdan MA. Supernumerary teeth: review of the literature and a survey of 152 cases. *Int J Paediatr Dent*. 2002 Jul;12(4):244–54.
30. Gawande P, Hebbale M, Patil T, Hindocha A, Halli R. Multiple Impacted Permanent and Supernumerary Teeth in the Anterior Mandible of Nonsyndromic Case: A Systematic Review and Multidisciplinary Approach to Management. *J Dent Allied Sci* 2015;4:47–51.
31. Ofer Sarne, Yehoshua Shapira, Sigalit Blumer, Tamar Finkelstein, Shirley Schonberger, Naomi Bechor, and Nir Shpack, Supernumerary Teeth in the Maxillary Anterior Region: The Dilemma of Early Versus Late Surgical Intervention. *Journal of Clinical Pediatric Dentistry*: 2018, Vol. 42, No. 1, pp. 55–61, 2018.
32. K. Khalaf, D. L. Robinson, C. Elcock, R. N. Smith, and A. H. Brook, "Tooth size in patients with supernumerary teeth and a control group measured by image analysis system," *Archives of Oral Biology*, vol. 50, no. 2, pp. 243–248, 2005.
33. Amaral D, Muthu M S. Supernumerary teeth: Review of literature and decision support system. *Indian J Dent Res* 2013;24:117–22.
34. Soames JV, Southam JC. *Textbook of oral pathology*. 2nd edn. Oxford, UK: Oxford University Press, 1993;. P6
35. C. L. Patchett, P. J. M. Crawford, A. C. Cameron, and C. D. Stephens, "The management of supernumerary teeth in childhood—a retrospective study of practice in Bristol Dental Hospital, England and Westmead Dental Hospital, Sydney, Australia," *International Journal of Paediatric Dentistry*, vol. 11, no. 4, pp. 259–265, 2001.
36. J. Foley, "Surgical removal of supernumerary teeth and the fate of incisor eruption," *European Journal of Paediatric Dentistry*, vol. 5, no. 1, pp. 35–40, 2004.
37. Kalaskar, R.R., Kalaskar, A.R. Multidisciplinary management of impacted central incisors due to supernumerary teeth and an associated dentigerous cyst. *Contemp Clin Dent*. 2011;2:53–58.
38. Shetty A, Hegde P, Shetty P, Udeshi P. Surgical management of supplemental tooth. *International Journal of Dentistry Research* 2017; 2(2): 26–28
39. Bahadure RN, ThosarN, JainES, Kharabe V, Gaikwad R. Supernumerary teeth in primary dentition and early intervention: Aseries of case reports. *Case Rep Dent* 2012;2012:614652.
40. Fazliah SN. Supernumerary tooth: report of a case. *Archives of Orofacial Sciences*. 2007 Oct;2:54–8.