Quantitative Assessment Of The Variations In Thyroid Dimensions In Relation To Increase In Serum TSH In Euthyroids

Mahrukh Kamran, Sahar Mubeen, Iffat Raza, Sanobar Bughio, Hira Waseem, Zainab Khalique Ansari

ABSTRACT:

Background: Volumetric evaluation of thyroid gland volume (TGV) is one of the criteria determining the iodine status of a population. TGV is the product of 3 dimension of each lobe: Anterio-posterior (AP) x medio-lateral (ML) x and cranio-caudal (CC) x correction factor.

Objectives: To determine the effect of serum TSH on thyroid dimensions of each lobe and to measure the amount of effect of per unit increase in serum TSH on thyroid dimensions of each lobe in euthyroids.

Study Design: It was a cross-sectional study.

Setting: The study was conducted at Ziauddin University Hospital, Clifton, Karachi.

Methodology: Healthy participants aged 21 years and above were included through convenient sampling. Serum Thyroid stimulating hormone was evaluated and ultrasound of thyroid gland TG of 192 euthyroid participants was performed. Spearman correlation and regression analysis was applied to evaluate the relationship between TSH and TG dimensions

Results: Relationship of increase in serum TSH with decrease in light lobe AP dimension was most significant. ($r = -0.142$ P-Value=0.001) and CC dimension least significant ($r = -0.0098$ P-Value=0.001). Where as in the left lobe AP dimension decreases significantly with increase serum TSH ($r = -0.147$ P-value=0.001). 11.7% of total variation in AP dimension, 3.5% of total variation in ML dimension and 6.5% of total variation in CC dimension in right lobe thyroid are because of serum TSH. While 9.5% of the total variation in AP dimension in left lobe is also due to serum TSH.

Conclusion: Negative and significant correlation between serum TSH and thyroid dimensions was observed. Serum TSH inversely and significantly affects all the dimensions of the right lobe and only one dimension in the left lobe.

KeyWords: Thyroid Gland, Thyrotropin, Diagnostic imaging

INTRODUCTION:

After urinary iodine level assessment, volumetric evaluation of thyroid gland (TG) has also become critical for determining the iodine status of a population as World Health Organization (WHO) suggested that normal range of thyroid gland volume (TGV) is required for large scale iodine monitoring programmes in different populations^{1,2}. Reference range of normal TG is important for diagnosing goiter, thyroiditis

and other thyroid diseases. TGV is also required to calculate the dose of radioiodine and also to evaluate the response of suppression treatment. It is also important to assess the TGV of such patients who are regularly undergoing long term treatment of the drugs that deranged TGV³. Determination of TGV is also vital for minimally invasive thyroid surgery⁴.

Ultrasonography of thyroid, due to its superficial location, is considered to be the best modality because it assures high sensitivity and specificity and good scanning speed^{5,7}. Due to its cost effectiveness and noninvasive approach it is considered to be the first choice to measure TGV and its morphology⁸. American Thyroid Association in 2009 suggested ultrasonography of thyroid as the most common and beneficial modality to image TG and its pathologies.

According to WHO recommended criteria to evaluate TGV, is to calculate the volume of each lobe of TG by the formula: Volume (ml) = Length in cm (Cranio-caudal dimension) x Width in cm (Medio-lateral dimension) x Depth in cm (Anterio-posterior dimension) x 0.4790. (0.479 = correction factor)^{1,2}. Total TGV is the sum of the volume of right and left thyroid lobes. Isthmus dimensions were not included in the measurement of thyroid gland volume as recommended by W.H.O^{1,2}.

Among the numbers of genetic and environmental factors that are known to effect TGV¹⁰⁻¹⁸, most extensively explored are the effect induced by iodine intake and thyroid stimulating

Mahrukh Kamran

Assistant Professor, DUHS, Ojha campus Karachi
Email: Mahrukhq1@hotmail.com

Sahar Mubeen

Associate Professor, DUHS, Ojha campus Karachi

Iffat Raza

Assistant Professor, Karachi Institute of Medical Science
Ojha campus Karachi

Sanobar Bughio

Senior Registrar, Department of radiology
Ziauddin University Hospital Clifton Karachi

Hira Waseem

Lecturer Department of Statistics, DUHS, Ojha campus Karachi

Zainab Khalique Ansari

Dow University Health Sciences
Ojha campus Karachi

Received: 16-05-2018

Revised: 24-05-2018

Accepted: 05-06-2018

hormone (TSH)^{12,14,19}. TSH and along with its receptor is important in, organization of the follicular cells of the thyroid and controlling TGV. After the binding of TSH to its receptors on the follicular cells, it then stimulates the growth of the follicular cells. These follicular cells in turn increase the synthesis and secretion of thyroxine²⁰⁻²².

There are studies that reported significantly negative correlation between serum TSH and TGV¹¹⁻¹³. Some of the studies also reported insignificant relationship between serum TSH and TGV^{10,23}. Volume of each lobe of thyroid gland is the product of its 3 dimension: Anterio-posterior(AP) x medio-lateral (ML) x cranio-caudal (CC) x correction factor^{1,2}. To the best of our knowledge no study reported the effect of serum TSH on thyroid dimensions. Objectives of this study is to determine the effect of serum TSH on thyroid dimensions in each lobe and also to measure the amount of effect of per unit increase in serum TSH on thyroid dimensions in each lobe.

Methodology:

A cross-sectional study was carried out at Ziauddin University hospital Karachi. This study was approved by ethical committee of Ziauddin University and hospitals. Apparently healthy subjects aged 21 years and above were recruited in the study through convenient sampling. Every participant was asked questions related to demographic profile, personal history, past history, medical and drug history. Participants with known thyroid diseases, with history of thyroid surgery, or on medications that are known to derange TGV (especially lithium, carbamazepine, phenytoin and oral contraceptive) were excluded from the study. Females with pregnancy, lactation and those who had delivered in last 12 months were also excluded due to risk of deranged TGV during this period.

Signs and symptoms related with thyroid functions were then noted to rule out unidentified cases of thyroid diseases. Physical examination of thyroid gland was then performed. All participants with enlarged thyroid gland and palpable thyroid nodule were excluded from the study. In order to recruit euthyroid participants, serum TSH level of the participants was then evaluated by chemiluminescence method. Reference range of TSH was 0.23 – 4.0 μ IU/ml as used by the laboratory of Ziauddin University Hospital.

Participants having Serum TSH between 0.23 – 4.0 μ IU/ml underwent ultrasonography of thyroid. Ultrasound machine Toshiba model SSA-590A with a 7.5 MHz transducer and 4 to 6 cm linear probe was used in this study. To avoid the inter observer error ultrasound was performed by a single radiologist. For the performance of ultrasound accurately standard technique for ultrasonography was followed, participants were examined in supine position, with pillow under their shoulders and their neck hyperextended. Ultrasound gel was applied all over the thyroid area of the neck. The probe was placed directly on the skin over the

thyroid gland. An image for each lobe of thyroid was obtained. Medio-lateral (ML) and antero-posterior (AP) dimensions of each lobe of thyroid were noted in transverse plane. Cranio-caudal (CC) dimension for each lobe of thyroid was noted in longitudinal plane^{1,2}.

Statistical Analysis:

Data was entered on Microsoft excel and analyzed on SPSS version 21. Mean with standard deviations were derived for serum TSH. Kolmogorov–Smirnov test was applied to check the normality. The data failed to follow the normality assumption therefore Log transformation was applied to normalize the data. Spearman correlation was applied for evaluating the correlation between thyroid dimensions and serum TSH level. Regression analysis was applied to evaluate the amount of variation in thyroid dimensions due to per unit increase in serum TSH. P-value < 0.05 was considered as significant.

RESULTS:

Mean serum TSH of 192 participants of the study was found to be 1.56 μ IU/ml $\pm$ 0.78 μ IU/ml with minimum and maximum 0.4 μ IU/ml to 3.74 μ IU/ml. Correlation Analysis between thyroid dimensions and serum TSH are shown in table 1. Change in thyroid dimension due to per unit increase in serum TSH shown in table 2. Graph1: Regression analysis determining the inverse linear relationship between serum TSH and Right Lobe Anterio-Posterior dimension of thyroid. (RT.AP LOG= Right lobe Anterio-Posterior Log transformation, TSHLOG=TSH Log transformation). Graph 2: Regression analysis determining the inverse linear relationship between serum TSH and Right Lobe Medio-Lateral dimension of thyroid. (RT.ML LOG= Right lobe Medio-Lateral Log transformation, TSHLOG=TSH Log transformation). Graph 3: Regression analysis determining the inverse linear relationship between serum TSH and Right Lobe Cranio-Caudal dimension of thyroid. (RT.CC LOG= Right lobe Cranio-Caudal Log transformation, TSHLOG=TSH Log transformation). Graph 4: Regression analysis determining the inverse linear relationship between serum TSH and Left Lobe Anterio-Posterior dimension of thyroid. (LT.AP LOG= Left lobe Anterio-Posterior Log transformation, TSHLOG=TSH Log transformation).

DISCUSSION:

Volumetric evaluation of TGV is one of the criteria determining the iodine status of a population. Volume of each lobe of TG is the product of its 3 dimension: Anterio-posterior(AP) x medio-lateral (ML) x cranio-caudal (CC) x correction factor^{1,2}. Previous studies reported either the inverse and significant relationship¹¹⁻¹³ or no relationship between serum TSH and TGV^{10,23}. To the best of our knowledge no study reported the effect of serum TSH on thyroid dimensions. Objectives of this study were to determine the effect of serum TSH on thyroid dimensions of each lobe and also to measure the amount of effect of per

N=192	Right Lobe Dimensions			Left Lobe Dimensions		
	Anterio-posterior	Medio-lateral	Cranio-caudal	Anterio-posterior	Medio-lateral	Cranio-caudal
Correlation	-0.321	-0.177	-0.280	-0.271	-0.127	-0.122
P-Value	0.001	0.014	0.001	0.0001	0.080	0.092

Table1: Correlation between serum TSH and Thyroid dimensions.

Correlation is significant at 0.05 level

N=192	Right Lobe Dimensions			Left Lobe Dimensions		
	Anterio-posterior	Medio-lateral	Cranio-caudal	Anterio-posterior	Medio-lateral	Cranio-caudal
Beta 1	-0.142	-0.1	-0.098	-0.147	-0.037	-0.039
B (constant)	0.174	0.223	0.449	0.158	0.200	0.434
R ²	0.117	0.035	0.065	0.095	0.004	0.012
P-Value	0.001	0.009	0.001	0.001	0.397	0.132

Table2: Regression analysis between thyroid dimension and serum TSH,

Correlation is significant at 0.05 level (2-tailed)

Graph 1: Regression between serum TSH and Right Lobe Anterio-Posterior Dimension R² Liner = 0.117 **Graph 2:** Regression between serum TSH and Right Lobe Medio-Lateral Dimension

unit increase in serum TSH on thyroid dimensions of each lobe.

Just like the inverse effect of serum TSH on TGV as reported previously^{11,13}, this study reported negative correlation between serum TSH and thyroid dimensions. In the right lobe of TG, AP dimension (P-Value 0.001), ML dimension (P-Value 0.014) and CC dimension (P-Value 0.001) dimension was found to be negatively and significantly related to serum TSH. Among the three dimensions of right lobe, serum TSH demonstrated most negative correlation with AP dimension. Least negative correlation was observed between serum TSH and ML dimension of right lobe.

Regression analysis further demonstrated that with per unit increase with serum TSH, AP dimension decreases significantly by 0.142ml, ML dimension decreases significantly by 0.1ml and CC dimension decreases by 0.098ml in the right lobe of thyroid. 11.7% of total variation in AP dimension, 3.5% of total variation in ML dimension and 6.5% of total variation in CC dimension of right lobe thyroid are the result of serum TSH.

In the left lobe of thyroid, significantly negative correlation was found only between serum TSH and AP dimension. Correlation between serum TSH and left lobe ML and CC dimension was found to insignificant. This study further

Graph 3: Regression between serum TSH and Right Lobe Cranio-Caudal Dimension

Graph 4: Regression between serum TSH and Left Lobe Antero-Posterior Dimension.

demonstrated that with per- unit increase in serum TSH only left lobe AP dimension decreased significantly by 0.147ml (P-Value=0.001). 9.5% of the total variation in AP dimension of left lobe was probably due to serum TSH. No correlation was reported between serum TSH and ML and CC dimension of left lobe of thyroid.

This study further demonstrated that if the effect of serum TSH was completely eliminated then in right lobe average of AP dimension would be 0.117cm, ML dimension would be 0.035 cm and CC dimension would be 0.06. In left lobe the average AP dimension would be 0.095 cm if the effect of serum TSH was removed.

Gomez et al., demonstrated negative and significant correlation between serum TSH and TGV in Spanish subjects ($r=-0.26$, P.value=0.001)¹³. Barrere et al also demonstrated inverse relationship between serum TSH and TGV among French subjects in both the genders¹¹. Negative correlation between the two variables was also observed in the Karachi population (P-Value=0.03) and also between serum TSH and right lobe volume (P-Value=0.029)¹².

The overall results of this study were not in accordance to the study done by Berghout *et al* in 1987 in Amsterdam population¹² Ressmussen et al in 1989 in Danish subjects²³ and Adibi et al in 2008 in Iranian population where they failed to report any relation between serum TSH and TGV and concluded that there was a lack of significant correlation between serum TSH level and TGV^{10,12,23}.

Conclusion:

Previous studies reported significant relationship between serum TSH and TGV or no relationship between serum TSH and TGV. To the best of our knowledge no study reported that which thyroid dimension in each lobe was most and least likely responsible for this negative correlation. Negative

and significant correlation between serum TSH and thyroid dimensions was observed in this study. Serum TSH inversely and significantly affected all the dimensions of the right lobe and only one dimension in the left lobe.

Acknowledgements: Ziauddin University and Hospitals.

Author Ship: This work was carried out in collaboration between all authors. Corresponding author Dr. Mahrugh Kamran designed the study, wrote the protocol, and wrote the first draft of the manuscript. Dr. Sahar Mubeen and Dr. Iffat Raza managed the literature search and critical analysis of the study. Author Dr. Sanobar Bughio supported the study technically and clinically. Hira Waseem and Dr. Zainab Khaiq Ansari participated in study design, data analysis and interpretation. All authors read and approved the final manuscript.

Conflict of interest: There was no conflict of interest.

REFERENCES:

1. Organization WH. Assessment of iodine deficiency disorders and monitoring their elimination: a guide for programme managers. 2007.
2. Organization WH, Disorders ICfCoID, UNICEF. Assessment of iodine deficiency disorders and monitoring their elimination: a guide for programme managers. 2001.
3. Dong BJ. Medicine Cabinet: How medications affect thyroid function. Western Journal of Medicine. 2000; 172(2): 102.
4. Hegazy MA, Khater AA, Setit AE, Amin MA, Kotb SZ, El Shafei MA, et al. Minimally invasive video-assisted thyroidectomy for small follicular thyroid nodules. World journal of surgery. 2007; 31(9): 1743-50.
5. Wiest PW, Hartshorne MF, Inskip PD, Crooks LA, Vela BS, Telepak RJ, et al. Thyroid palpation versus high-resolution thyroid ultrasonography in the detection of nodules. Journal of Ultrasound in Medicine. 1998; 17(8): 487-96.

6. Kangelaris GT, Kim TB, Orloff LA. Role of ultrasound in thyroid disorders. *Otolaryngologic Clinics of North America*. 2010; 43(6): 1209-27.
7. Zimmermann M, Saad A, Hess S, Torresani T, Chaouki N. Thyroid ultrasound compared with World Health Organization 1960 and 1994 palpation criteria for determination of goiter prevalence in regions of mild and severe iodine deficiency. *European Journal of Endocrinology*. 2000; 143(6): 727-31.
8. Hegedüs L. Thyroid ultrasound. *Endocrinology and Metabolism Clinics*. 2001; 30(2): 339-60.
9. Cooper DS, Doherty GM, Haugen BR, Kloos RT, Lee SL, Mandel SJ, et al. Revised American Thyroid Association management guidelines for patients with thyroid nodules and differentiated thyroid cancer: the American Thyroid Association (ATA) guidelines taskforce on thyroid nodules and differentiated thyroid cancer. *Thyroid*. 2009; 19(11): 1167-214.
10. Adibi A, Sirous M, Aminorroaya A, Roohi E, Mostafavi M, Fallah Z, et al. Normal values of thyroid gland in Isfahan, an iodine replete area. *Journal of Research in Medical Sciences*. 2008; 13(2): 55-60.
11. Barrère X, Valeix P, Preziosi P, Bensimon M, Pelletier B, Galan P, et al. Determinants of thyroid volume in healthy French adults participating in the SU. VI. MAX cohort. *Clinical endocrinology*. 2000; 52(3): 273-8.
12. Mahrukh Kamran SM, Iffat Raza, Sanobar Bughio, Hira Waseen, Zainab Khaliq Ansari. The relationship of thyroid gland volume (TGV) with thyroid stimulating hormone (TSH) and the quantitative measurement of the variations in TGV in relation to per unit increase in serum tsh. *European Journal of Biomedical and Pharmaceutical Sciences*. 2018; 5(5): 1154-8.
13. Gómez JM, Maravall FJ, Gómez N, Gumà A, Soler J. Determinants of thyroid volume as measured by ultrasonography in healthy adults randomly selected. *Clinical endocrinology*. 2000; 53(5): 629-34.
14. Hegedüs L. Thyroid size determined by ultrasound. Influence of physiological factors and non-thyroidal disease. *Danish medical bulletin*. 1990; 37(3): 249-63.
15. Turcios S, Lence-Anta JJ, Santana J-L, Pereda CM, Velasco M, Chappe M, et al. Thyroid volume and its relation to anthropometric measures in a healthy Cuban population. *European thyroid journal*. 2015; 4(1): 55-61.
16. HEGEDÜS L, Perrild H, Poulsen LR, ANDERSEN JR, Holm B, SCHNOHR P, et al. The determination of thyroid volume by ultrasound and its relationship to body weight, age, and sex in normal subjects. *The Journal of Clinical Endocrinology & Metabolism*. 1983; 56(2): 260-3.
17. Moghadam RN, Shajari A, Afkhami-Ardekani M. Influence of physiological factors on thyroid size determined by ultrasound. *Acta Medica Iranica*. 2011; 49(5): 302.
18. Ivanac G, Rožman B, Škreb F, Brkljač B, Pavić L. Ultrasonographic measurement of the thyroid volume. *Collegium antropologicum*. 2004; 28(1): 287-91.
19. Vejbjerg P, Knudsen N, Perrild H, Carlé A, Laurberg P, Pedersen IB, et al. Effect of a mandatory iodization program on thyroid gland volume based on individuals' age, gender, and preceding severity of dietary iodine deficiency: a prospective, population-based study. *The Journal of Clinical Endocrinology & Metabolism*. 2007; 92(4): 1397-401.
20. Dumont JE, Lamy F, Roger P, Maenhaut C. Physiological and pathological regulation of thyroid cell proliferation and differentiation by thyrotropin and other factors. *Physiological reviews*. 1992; 72(3): 667-97.
21. De Felice M, Postiglione MP, Di Lauro R. Minireview: thyrotropin receptor signaling in development and differentiation of the thyroid gland: insights from mouse models and human diseases. *Endocrinology*. 2004; 145(9): 4062-7.
22. Postiglione M, Parlato R, Rodriguez-Mallon A, Rosica A, Mithbaokar P, Maresca M, et al. Role of the thyroid-stimulating hormone receptor signaling in development and differentiation of the thyroid gland. *Proceedings of the National Academy of Sciences*. 2002; 99(24): 15462-7.
23. Feldt-Rasmussen U, Hegedüs L, Perrild H, Rasmussen N, Hansen J. Relationship between serum thyroglobulin, thyroid volume and serum TSH in healthy non-goitrous subjects and the relationship to seasonal variations in iodine intake. *Thyroidology*. 1989; 1(3): 115-8.